

Technical Fact Sheet – N-Nitroso-dimethylamine (NDMA)

January 2017

TECHNICAL FACT SHEET- NDWA

At a Glance

- Formerly used in the production of rocket fuel, antioxidants and softeners for copolymers.
 Currently used only for research purposes.
- Unintended byproduct of chlorination of wastewater at wastewater treatment plants that use chloramines for disinfection, raising significant concern as a drinking water contaminant.
- Highly mobile in soil, with potential to leach into groundwater.
- ☐ Oral route is the primary human exposure pathway.
- Classified as a B2 (probable human) carcinogen.
- □ Listed as a priority pollutant by the EPA, but no federal standard has been established for drinking water.
- Detection methods include solid phase extraction, gas chromatography and liquid chromatography.
- Most common NDMA water cleanup method is via photolysis by ultraviolet radiation. Potential for aerobic and anaerobic NDMA biodegradation also exists.

Introduction

This fact sheet, developed by the U.S. Environmental Protection Agency (EPA) Federal Facilities Restoration and Reuse Office (FFRRO), provides a summary of the contaminant N-Nitrosodimethylamine (NDMA), including physical and chemical properties; environmental and health impacts; existing federal and state guidelines; detection and treatment methods; and additional sources of information. This fact sheet is intended for use by site managers and other field personnel who may address NDMA contamination at cleanup sites or in drinking water supplies.

NDMA is a drinking water contaminant of concern because of its miscibility with water, as well as its carcinogenicity and toxicity.

What is NDMA?

- □ NDMA is a semivolatile organic chemical that forms in both industrial and natural processes (Cal/EPA 2006; Mitch and others 2003b).
- □ NDMA is not currently produced in pure form or commercially used in the United States, except for research purposes. It was formerly used in production of liquid rocket fuel, antioxidants, additives for lubricants and softeners for copolymers (ATSDR 1989; HSDB 2013).
- □ NDMA can be unintentionally produced in and released from industrial sources through chemical reactions, such as those that involve alkylamines. Potential industrial sources include amine manufacturing plants, tanneries, pesticide manufacturing plants, rubber and tire manufacturers, fish processing facilities, foundries, dye manufacturers and surfactant industries (ATSDR 1989).
- □ NDMA is also an unintended byproduct of the chlorination of wastewater and drinking water at treatment plants that use chloramines for disinfection (Bradley and others 2005; Mitch and others 2003).

<u>Disclaimer:</u> The U.S. EPA prepared this fact sheet from publicly-available sources; additional information can be obtained from the source documents. This fact sheet is not intended to be used as a primary source of information and is not intended, nor can it be relied upon, to create any rights enforceable by any party in litigation with the United States. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

Exhibit 1: Physical and Chemical Properties of NDMA (ATSDR 1989; Cal/EPA 2006; HSDB 2013; NIOSH 2016)

Property	Value/Description
Chemical Abstract Systems (CAS) Number	62-75-9
Physical description (physical state at room temperature)	Yellow liquid with faint or no odor
Molecular weight (g/mol)	74.08
Water solubility at 25°C	Miscible
Melting point (°C)	-25 (estimated)
Boiling point (°C)	152 to 154
Specific gravity/Density at 20 °F/4 °C (g/mL)	1.005 to 1.006
Vapor pressure at 20°C (mm Hg)	2.7
Organic carbon partition coefficient (log K _∞)	1.07 (estimated)
Octanol-water partition coefficient (log Kow)	-0.57
Henry's Law Constant at 20°C (atm-m³/mol)	2.63 x 10 ⁻⁷ (ATSDR 1989) 1.08 x 10 ⁻⁶ (HSDB 2013)

Abbreviations: g/mol – grams per mole; °C – degrees Celsius; g/mL – grams per milliliter; mm Hg – millimeters of mercury; atm-m³/mol – atmosphere-cubic meters per mole.

What are the environmental impacts of NDMA?

In a 2002 survey conducted by the California NDMA contamination may be found in air, soil and Department of Health Services (CDHS), elevated water (ATSDR 1999). concentrations of NDMA were detected in When released to the air, NDMA is expected to locations where wastewater treatment plant exist solely as vapor in the ambient atmosphere effluent was used for aquifer recharge and near and is broken down quickly by sunlight within facilities that use unsymmetrical dimethylhydrazine minutes (HSDB 2013). (UDMH)-based rocket fuel (CDHS 2002; Mitch and When released to soil, NDMA can be highly others 2003b). mobile and will either volatilize or leach into As of March 2011, NDMA was the predominant groundwater (ATSDR 1999; HSDB 2013). nitrosamine detected in samples obtained from In water, NDMA is completely miscible and is not public water systems, which were monitored as expected to sorb onto solid particles or sediment. part of the unregulated contaminant monitoring NDMA may break down in water as a result of rule (UCMR). The EPA uses the UCMR to monitor exposure to sunlight or by natural biological contaminants that are suspected to be present in processes. The potential for bioconcentration in drinking water but that do not currently have aquatic organisms is low based on an estimated health-based standards under the Safe Drinking bioconcentration factor of 3 (ATSDR 1999; HSDB Water Act (EPA 2011a; EPA 2014). 2013; WHO 2008). The second UCMR was analyzed for NDMA At rocket engine testing facilities in California, occurrence and trends across the U.S. NDMA NDMA has been found at high concentrations in occurrence was strongly associated with groundwater on site (up to 400,000 nanograms chloramine use. Elevated NDMA was more per liter [ng/L]) and also in downgradient drinking common in surface water systems than water wells (up to 20,000 ng/L) (Mitch and others groundwater systems. Smaller utilities were found 2003b). to have the most extreme NDMA levels (Woods and Dickenson 2015)

What are the routes of exposure and health effects of NDMA?

NDMA exposure may occur through (1) ingesting function of liver, kidneys and lungs; and dizziness food that contains nitrosamines, such as smoked (HSDB 2013; OSHA 2005). or cured meats and fish; (2) ingesting food that EPA has classified NDMA as a B2 (probable contains alkylamines, which can cause NDMA to human) carcinogen based on the induction of form in the stomach; (3) drinking contaminated tumors at multiple sites in different mammal water; (4) drinking malt beverages (such as beer species exposed to NDMA by various routes (EPA and whiskey) that may contain low levels of IRIS 2002). nitrosamines formed during processing; (5) using The U.S. Department of Health and Human toiletry and cosmetic products such as shampoos Services (DHHS) states that NDMA is reasonably and cleansers that contain NDMA; and (6) anticipated to be a human carcinogen (NTP 2014). breathing or inhaling cigarette smoke. Workplace DHHS states that NDMA caused tumors in exposure can occur at tanneries, pesticide numerous species of experimental animals, at manufacturing plants and rubber and tire plants several different tissue sites, and by several (ATSDR 1989, 1999). different routes of exposure. Tumors occurred The oral route, including consumption of primarily of the liver, respiratory tract, kidney and contaminated food and water, is the primary blood vessels (NTP 2014; IARC 1998). human exposure pathway for NDMA (ATSDR The American Conference of Governmental 1989; Cal/EPA 2006). Industrial Hygienists (ACGIH) has classified Exposure to high levels of NDMA may cause liver NDMA as a Group A3 confirmed animal damage in humans (ATSDR 1999; HSDB 2013). carcinogen with unknown relevance to humans Potential symptoms of overexposure include (HSDB 2013). headache; fever; nausea; jaundice; vomiting; abdominal cramps; enlarged liver; reduced

Are there any federal and state guidelines and health standards for NDMA?

EPA has not derived a chronic oral reference dose (RfD) or a chronic inhalation reference concentration (RfC) for evaluating NDMA's noncancer effects in the EPA's Integrated Risk Information System database (EPA IRIS 2002).
 EPA has derived a RfD of 8.0 x 10 ⁻⁶ mg/kg-day and an RfC of 4.0 x 10 ⁻⁵ mg/m ³ as Provisional Peer-Reviewed Toxicity Values (PPRTVs) for evaluating noncancer effects (EPA 2007).
 EPA has assigned an oral slope factor for carcinogenic risk of 51 milligrams per kilogram per day (mg/kg-day) ⁻¹ , a drinking water unit risk of 1.4 x 10^{-3} per microgram per liter (μ g/L) ⁻¹ and an inhalation unit risk of 1.4 x 10^{-2} μ g per cubic meter (m³) (EPA IRIS 2002).
For tap water, EPA calculated a screening level of 0.11 ng/L for NDMA, based on a 10 ⁻⁶ lifetime excess cancer risk (EPA 2016).
 EPA's screening levels for soil are 2.0×10^{-3} milligrams per kilogram (mg/kg) for residential and 3.4×10^{-2} mg/kg for industrial (based on 10^{-6} cancer risk). The soil screening level for protection of groundwater is 2.7×10^{-8} mg/kg (EPA 2016).
EPA's screening levels for air are 7.2×10^{-5} micrograms per cubic meter (μ g/m³) for residential and $8.8 \times 10^{-4} \mu$ g/m³ for industrial (based on 10^{-6}

cancer risk) (EPA 2016).

Various states have established drinking water and groundwater guidelines, including the following:

State	Guideline (µg/L)	Source
Alabama	0.0013	ADEM 2008
Alaska	0.017	AL DEC 2008
California	0.003	Cal/EPA 2006
Colorado	0.00069	CDPHE 2013
Delaware	0.001	DE DNR 1999
Florida	0.0007	FDEP 2005
Indiana	0.0049	IDEM 2015
Massachusetts	0.01	MADEP 2004
Mississippi	0.00131	MS DEQ 2002
New Jersey	0.0007	NJDEP 2015
North Carolina	0.0007	NCDENR 2015
Pennsylvania	0.0014	PADEP 2011
Texas	0.018	TCEQ 2016
Washington	0.000858	WA DEP 2015
West Virginia	0.0013	WV DEP 2009

Technical Fact Sheet - NDMA

Are there any federal and state guidelines and health standards for NDMA? (continued)

1	Onthina Caj		
	EPA included NDMA on the third Contaminant Candidate List (CCL3), which is a list of unregulated contaminants that are known to or anticipated to occur in public water systems and may require regulation under the Safe Drinking Water Act (EPA 2009; EPA 2011a; EPA 2014).		In addition, EPA added NDMA to its UCMR 2, requiring many large water utilities to monitor for NDMA (EPA 2015).
N	hat detection and site characterization	on	methods are available for NDMA

□ For drinking water, EPA Method 521 uses solid phase extraction (SPE) and capillary column gas chromatography (GC) with large-volume as alkyl injection and chemical ionization tandem mass spectroscopy (MS) (EPA 2004). □ For wastewater, EPA Method 607 uses methylene chloride extraction, GC and a

- For wastewater, EPA Method 607 uses methylene chloride extraction, GC and a nitrogen-phosphorus detector (NPD) (EPA 2007; EPA 2016).
- ☐ For wastewater, EPA Method 1625 uses isotope dilution, GC and MS (EPA 2007; EPA 2016).
- □ For groundwater, wastewater, soil, sediment and sludges, EPA SW-846 Method 8070 uses methylene chloride extraction, GC and a NPD (EPA 1996).
- For solid waste matrices, soil, air sampling media and water samples, EPA SW-846 Method 8270 uses GC and MS (EPA 1998).

- An analytical method has also been developed specifically to quantify NDMA precursors such as alkylamines in waste or wastewater (Mitch, and others 2003).
- A method using liquid chromatography tandem MS (LC/MS/MS) detects both thermally stable and unstable nitrosamines in drinking water (Zhao and others 2006).
- □ A study developed a method that is a combination of SPE and LC/MS/MS for determination of NDMA in surface water, groundwater and wastewater samples. The quantification limit identified was 2 ng/L (Topuz and others 2012).
- Modifications to GC-MS and GC-NPD methods including sample evapoconcentration and low concentration instrument calibration can be used to detect NDMA in soil to levels below 1 microgram per kilogram (μg/kg) (USACE 2009).

What technologies are being used to treat NDMA?

- ☐ The most common method to treat NDMA in drinking water systems is photolysis by ultraviolet radiation in the wavelength range of 225 to 250 nanometers (nm) (Mitch and others 2003b).
- Biological treatment, microfiltration and reverse osmosis treatment may be used to remove NDMA precursors from wastewater before chlorination (Mitch and others 2003b).
- Activated sludge, biological activated carbon and ultraviolet photolysis were found to be effective for NDMA mitigation in a study investigating 11 sites using ozone-based wastewater treatment trains (Gerrity and others 2015).
- ☐ The Department of Defense's Strategic Environmental Research and Development Program (SERDP) is investigating abiotic, biotic and coupled abiotic/biotic processes to accelerate NDMA degradation in the subsurface (DoD SERDP 2008, 2009, 2012).

- A recent study of NDMA precursors found that photolysis and biodegradation were effective removal mechanisms for precursors in the water column (Woods and Dickenson 2016).
- Laboratory-scale studies have shown that aerobic and anaerobic biodegradation of NDMA to low ng/L concentrations in water and soil may be possible (Bradley and others 2005; DoD SERDP 2008).
- A laboratory-scale study demonstrated the potential for in-situ aerobic cometabolism of NDMA in the presence of methane- and benzene-amended groundwater highlighting possible attenuation mechanisms and rates for NDMA biotransformation in aerobic aquifers undergoing active remediation, natural attenuation or managed aquifer recharge with treated wastewater (Weidhaas and Dupont 2013).

What technologies are being used to treat NDMA (continued)?

An Environmental Security Technology involved in the aerobic biodegradation of NDMA. Certification Program demonstration project Laboratory-scale study results highlighted the evaluated the technical effectiveness and cost of importance of monooxygenases in the degradation using a fluidized bed bioreactor (FBR) for treating of NDMA (DoD SERDP 2012). NDMA in groundwater at a test facility. The FBR A SERDP field study was recently completed was found to be an effective means to treat utilizing propane biosparging for in situ NDMA, decreasing concentrations from 1 µg/L to remediation of NDMA in groundwater. The field 4.2 ng/L. The cost of the full-scale FBR was test results support that propane biosparging can determined to be significantly less than the be an effective approach to reduce the comparable ultraviolet system over a 30-year concentrations of NDMA in a groundwater aguifer remedial timeframe (ESTCP 2014). by 3 to 4 orders of magnitude, and that Laboratory-scale study results suggest that in-situ concentrations in the low nanograms per liter coupled abiotic/biotic processes may efficiently (ng/L) range can be achieved with continuous degrade NDMA in groundwater (DoD SERDP treatment (DoD SERDP 2016). 2009). A laboratory-scale study observed the Membrane bioreactor (MBR) treatment was found decomposition of NDMA in water using nanoscale to be effective in removing NDMA through zero-valent iron in the presence of aluminum and biodegradation due to the presence of strong iron salts. The highest removal was found at a pH electron donating functional groups in their of 5. Improved removal was associated with a structure (Wijekoon and others 2013). higher reaction temperature (Lin Lin and others

2013).

Where can I find more information about NDMA?

An SERDP project was conducted to identify the organisms, enzymes and biochemical pathways

2010. "Drinking Water Notification Levels and

	Registry (ATSDR). 1989. "Toxicological Profile for <i>n</i> -Nitrosodimethylamine."	www.cdph.ca.gov/certlic/drinkingwater/Documents/Notificationlevels/notificationlevels.pdf
	www.atsdr.cdc.gov/toxprofiles/tp141.pdf Alabama Department of Environmental Management (ADEM). 2008. "Alabama Risk- Based Corrective Action Guidance Manual." http://adem.alabama.gov/programs/land/landforms/ARBCAManual.pdf	California Environmental Protection Agency (Cal/EPA). Office of Environmental Health Hazard Assessment. 2006. "Public Health Goals for Chemicals in Drinking Water - N-Nitrosodimethylamine." http://oehha.ca.gov/media/downloads/water/chemi
	Alaska Department of Environmental Conservation (AL DEC) Division of Water. 2008. Groundwater Cleanup Levels. http://dec.alaska.gov/SPAR/csp/guidance_forms/docs/Groundwater_Cleanup_Levels.pdf ATSDR. 1999. "ToxFAQs - N-	cals/phg/122206ndmaphg_0.pdf Colorado Department of Public Health and Environment (CDPHE) Water Quality Control Commission. 2013. "The Basic Standards for Groundwater." https://www.colorado.gov/pacific/sites/default/files/
	Nitrosodimethylamine." https://www.atsdr.cdc.gov/toxfaqs/tfacts141.pdf Bradley, P.M., Carr, S.A., Baird, R.B., and F.H. Chappelle. 2005. "Biodegradation of N-	 41_2016%2806%29hdr.pdf Delaware Department of Natural Resources and Environmental Control (DE DNREC). 1999. "Remediation Standards Guidance."
	nitrosodimethylamine in Soil from a Water Reclamation Facility." Bioremediation Journal. Volume 9. Pages 115 to 120. https://www.researchgate.net/publication/2489503 13 Biodegradation of Nnitrosodimethylamine in Soil from a Water Reclamation Facility	http://www.dnrec.state.de.us/DNREC2000/Divisions/AWM/sirb/DOCS/PDFS/Misc/RemStnd.pdf Florida Department of Environmental Protection (FDEP). 2005. "Contaminant Cleanup Target Levels." https://www.flrules.org/gateway/ChapterHome.asp?Chapter=62-777
Ш	California Department of Public Health (CDPH).	

Where can I find more information about NDMA? (continued)

Gerrity, D., Pisarenko, A.N., Marti, E., Trenholm, R.A., Gerringer, F., Reungoat, J. and Dickenson, E. 2015. "Nitrosamines in pilot-scale and full-scale	RC/ERC%20Reports%20Received/2015/Departm ent%20of%20Environment%20and%20Natural%2 0Resources/2015-
wastewater treatment plants with ozonation." Water Research. Volume 72. Pages 251 to 261.	Jan%20Study%20Use%20of%20Contaminated%20Property.pdf
Hazardous Substance Data Bank (HSDB). 2013. N-Nitrosodimethylamine. http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB	Occupational Safety and Health Administration (OSHA). 2005. Chemical Sampling Information – N-Nitrosodimethylamine. www.osha.gov/dts/
 International Agency for Research on Cancer (IARC). 1998. "N-Nitrosodimethylamine." Some N-nitroso compounds. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Volume 17. Lyon, France: International Agency for Research on Cancer. Page 125. http://www.inchem.org/documents/iarc/vol17/n-nitrosodimethylamine.html	chemicalsampling/data/CH_258000.html. Pennsylvania Department of Environmental Protection (PADEP). 2011. Statewide Health Standards. http://files.dep.state.pa.us/EnvironmentalCleanupErownfields/LandRecyclingProgram/LandRecyclingProgramPortalFiles/SWHTables/Table%201%202011.pdf
Indiana Department of Environmental Management (IDEM). 2015. "Remediation Closure Guide." http://www.in.gov/idem/cleanups/files/remediation closure guide.pdf	 Texas Commission on Environmental Quality (TCEQ). 2016. "Texas Risk Reduction Program Rule." http://www.tceq.texas.gov/assets/public/remediation/trrp/pcls.xlsx
 Lin Lin, B. X., Lin, Y., Yan, L., Shen, K., Xia, S., Hu, C. and Rong, R. 2013. "Reduction of <i>N</i> -Nitrosodimethylamine (NDMA) in Aqueous Solution by Nanoscale Fe/Al2(SO4)3." Water, Air, & Soil Pollution. Volume 224 (7). Page 1.	Topuz, E., Aydin, E., and E. Pehlivanoglu-Mantas. 2012. "A Practical LC-MS/MS Method for the Detection of NDMA at Nanogram per Liter Concentrations in Multiple Water Matrices." Water, Air, & Soil Pollution. Volume 223 (9). Pages 5793 to 5802.
Massachusetts Department of Environmental Protection (Mass DEP). 2004. "Current Regulatory Limit: n-Nitrosodimethylamine." www.mass.gov/eea/agencies/massdep/water/drinking/standards/n-nitrosodimethylamine-ndma.html	U.S. Army Corps of Engineers (USACE). 2009. "Determination of Low Level NDMA in Soils." ERDC TN-EQT-09-01. http://acwc.sdp.sirsi.net/client/en_US/default/index_assetbox.assetactionicon.view/1045326?rm=ENV
Mississippi Department of Environmental Quality (MS DEQ). 2002. "Risk Evaluation Procedures for Voluntary Cleanup and Redevelopment of	IRONMENTAL+1%7C%7C%7C1%7C%7C%7C0 %7C%7C%7Ctrueℑ=WES
Brownfield Sites." http://www.deq.state.ms.us/MDEQ.nsf/pdf/GARD brownfieldrisk/\$File/Proced.pdf?OpenElement	 U.S. Department of Defense (DoD) Strategic Environmental Research and Development Program (SERDP). 2008. "Bioremediation Approaches for Treating Low Concentrations of
Mitch, W.A., Sharp, J.O, Trussell, R.R., Valentine, R.L, Alvarez-Cohen, L., and D.L. Sedlack. 2003b. "N-Nitrosodimethylamine (NDMA) as a Drinking Water Contaminant: A Review." Environmental	NNitrosodimethylamine in Groundwater." SERDP Project ER-1456. http://www.dtic.mil/cgi-bin/GetTRDoc?Location=U2&doc=GetTRDoc.pdf8 https://www.dtic.mil/cgi-bin/GetTRDoc.pdf8 https://www.dtic.mil/cgi-bin/GetTRDoc.pdf8 https://www.dtoc.mil/cgi-bin/GetTRDoc.pdf8 <a 1421<="" abiotic="" and="" biotic="" controlling="" ep="" href="https://www.dtoc.mil/cgi-</td></tr><tr><td>Engineering Science. Volume 20 (5). Pages 389 to 404. http://superfund.berkeley.edu/pdf/231.pdf</td><td>DoD SERDP. 2009. " in="" mechanisms="" ndma."="" of="" project="" remediation="" serdp="" situ="" td="">
New Jersey Department of Environmental Protection (NJDEP). 2015. "Ground Water Quality Standards – Class IIA by Constituent." http://www.nj.gov/dep/standards/ground%20water.pdf	NDMA." SERDP Project ER-1421. https://docs.serdp- estcp.org/content/download/6404/85420/file/ER- 1421 Final Report.pdf
North Carolina Department of Environment and Natural Resources(NCDENR). 2015. "Study Use of Contaminated Property, Risk Based Report."	DoD SERDP 2012. "Oxygenase-Catalyzed Biodegradation of Emerging Water Contaminants: 1,4-Dioxane and N-Nitrosodimethylamine." SERDP Project ER-propane. https://docs.serdp-

http://www.ncleg.net/documentsites/committees/E

<u>estcp.org/content/download/15286/174933/file/ER-1417-FR.pdf</u>

Where can I find more information about NDMA? (continued)


		,
DoD SERDP 2016. "Field Demonstration of Propane Biosparging for In Situ Remediation of N-Nitrosodimethylamine (NDMA) in Groundwater."		EPA. 1998. "Methods 8270D. Semivolatile Organic Compounds by Gas Chromatography/Mass Spectrometry (GC/MS)."
SERDP Project ER-200828. https://www.serdp-estcp.org/content/download/40059/384461/file/Final%20Report.V1%20ER-		https://www.epa.gov/sites/production/files/2015- 07/documents/epa-8270d.pdf
200828%20January%202016.pdf Environmental Security Technology Certification Program (ESTCP). 2014. "Treatment of N-		EPA. 2009. "Drinking Water Contaminant Candidate List 3 – Final." Federal Register Notice. https://www.gpo.gov/fdsys/pkg/FR-2009-10-
Nitrosodimethylamine in Groundwater Using a Fluidized Bed Bioreactor". https://docs.serdp-estcp.org/content/download/24415/252901/file/ER-200829-FR.pdf		 <u>08/pdf/E9-24287.pdf</u> EPA. 2011a. "Regulatory Determinations for the Third Drinking Water Contaminant Candidate List." Stakeholder Meeting. June 6, 2011. Washington
 National Toxicology Program (NTP). 2014. Report on Carcinogens, Thirteenth Edition. Research Triangle Park, NC: U.S. Department of Health and Human Services, Public Health Service. https://www.niehs.nih.gov/health/material		D.C. https://www.epa.gov/sites/production/files/2014-09/documents/preliminary-regulatory-determinations-3-june-16th-public-meeting-slides.pdf
s/report on carcinogens 13th edition the 508.p	Ц	EPA. 2014. "Preliminary Regulatory Determinations for the Third Drinking Water
U.S. Environmental Protection Agency (EPA).1996. "Method 8070A. Nitrosamines By Gas Chromatography."		Contaminant Candidate List." Stakeholder Meeting. December 9, 2014. Washington D.C. https://www.epa.gov/sites/production/files/2015-
https://www.epa.gov/sites/production/files/2015- 12/documents/8070a.pdf		07/documents/preregdet3stakeholderbriefingdec2 014.pdf
EPA. 2007. Provisional Peer Reviewed Toxicity Values for N-Nitrosodimethylamine (CASRN 62-75-9). 6-19-2007. https://hhpprtv.ornl.gov/issue_papers/NitrosodimethylamineN.pdf		EPA. 2013. "Toxics Criteria for those States Not Complying with Clean Water Act Section 303(c)(2)(B)." Code of Federal Regulations (CFR). Title 40, Chapter 1, Part 131.36. July 1 edition. https://www.gpo.gov/fdsys/pkg/CFR-2014-title40-vol22-sec131-36.pdf
EPA. 2010. "Designation of a Hazardous Substance." Code of Federal Regulations (CFR). Title 40, Chapter 1, Part 302.4.	Ц	EPA. 2015. Second Unregulated Contaminant Monitoring Rule 2 (UCMR 2).
EPA. 2004. "U.S. EPA Method 521: Determination of Nitrosoamines in Drinking Water		https://www.epa.gov/dwucmr/second-unregulated-contaminant-monitoring-rule
by Solid Phase Extraction (SPE) and Capillary Column Gas Chromatography with Large Volume Injection and Chemical Ionization Tandem Mass		EPA. 2016. Approved Clean Water Act Test Methods: Organic Chemical Analysis. water.epa.gov/scitech/methods/cwa/organics
Spectrometry (MS/MS)." Version 1.0. National Exposure Research Laboratory, Cincinnati, Ohio. EPA 600-R-05-054.		EPA. 2016. Regional Screening Level (RSL) Summary Table. https://www.epa.gov/risk/regional-screening-
https://cfpub.epa.gov/si/si public file download.cf m?p download id=525080		levels-rsls-generic-tables-may-2016 EPA Integrated Risk Information System (IRIS).
EPA. 2007. "Guidelines Establishing Test Procedures for the Analysis of Pollutants Under		2002. "N-Nitrosodimethylamine; CASRN 62-75-9." https://cfpub.epa.gov/ncea/iris/iris_documents/doc uments/subst/0045_summary.pdf.
the Clean Water Act; National Primary Drinking Water Regulations; and National Secondary Drinking Water Regulations; Analysis and Sampling Procedures; Final Rule." 40 CFR Part 122, 136, et seq.		Washington Department of Ecology (Ecology). 2015. "Groundwater Methods B and A ARARs." https://fortress.wa.gov/ecy/clarc/FocusSheets/Groundwater%20Methods%20B%20and%20A%20and%20ARARs.pdf

Technical Fact Sheet - NDMA

Webster, T.S., Condee, C., and P.B. Hatzinger.
 2013. "Ex situ treatment of N-nitrosodimethylamine (NDMA) in Groundwater

using a Fluidized Bed Reactor." Water Research. Volume 47 (2). Pages 811 to 820.

Where can I find more information about NDMA? (continued)



Contact Information

If you have any questions or comments on this fact sheet, please contact: Mary Cooke, FFRRO, by email at cooke.maryt@epa.gov.